Geology of the Lafayette 7.5-Minute Quadrangle, LA

Louisiana Geological Survey

Introduction, Location, and Geologic Setting

The study area lies within the Gulf Coast salt basin, directly north of the northwestern terminus of the Five Islands salt-dome trend in southwestern Louisiana. It lies directly west of the western valley wall of the Holocene Mississippi River flood plain near the southern edge of coast-parallel outcrop belts of terraced Pleistocene strata.

The area covers portions of two parishes (Figures 1, 2), Lafayette and Vermilion. The basic framework of surface geology of the region encompassing the study area was detailed by Howe and Moresi (1931, 1933) and Howe et al. (1935), and later was rendered at approximately 1:1,056,000 scale by Jones et al. (1954, Plate I) drawing in part upon unpublished work by Fisk (1948). Busch et al. (1974) differentiated Holocene Mississippi floodplain alluvium in the Atchafalaya basin, including the area occupied by the Mississippi River flood plain to the east of the Lafayette quadrangle. Loess flanking the lower Mississippi River valley in Louisiana was investigated and mapped by Miller (1983) and summarized by Miller et al. (1985). Reviews and mapping of local environmental geology that include the proposed study area were prepared by Rouly (1989) and Saxton (1986). The regional framework and context of the study area was updated at 1:1,100,000 scale in the compilation by Saucier and Snead (1989). Autin et al. (1991) and Saucier (1994a) reviewed, updated, and summarized the surface geology of the lower Mississippi valley. Saucier (1994b, his plate 11) revised the mapping of Mississippi River floodplain alluvium at 1:250,000 scale. Aspects of sediments of the Avoyelles Prairie (Avoyelles alloformation)compilation 2 Tm[(37-19(y)10(e)4(ll)-3(e)4(s)-110(a)4(ll)-3(ot

aquifer; and Van Biersel and Milner (2010) summarized the aquifer's distribution, recharge area, proportions of water-use categories, and pumpage rates.

Methods



2. Surface geology of the greater Lafayette area and environs (mosaicked excerpts adapted from Heinrich et al., 2003, and Heinrich and Autin, 2000). (**Ppbe**, Beaumont Alloformation; **Ppav**, Avoyelles alloformation; **Ppbc**, Big Cane alloformation; **Hmd$_1$**, Distributary complex of Mississippi River meander belt 1; **Hmc$_3$u**, Crevasse complex of Mississippi River meander belt 3, upper deposits; **Hmd$_3$u**, Distributary complex of Mississippi River meander belt 3, upper deposits; **Hml$_3$u**, Natural levee complex of Mississippi River meander belt 3, upper deposits; **Hmm$_3$l**, Mississippi River meander belt 3, lower deposits; **Hb**, Backswamp deposits; **Hrm**, Red River meander-belt deposits; **Hala**, Lacustrine deposits associated with the Atchafalaya River; **Hal**, Atchafalaya River natural levee deposits; **Hac**, Atchafalaya River crevasse splay deposits; **Had**, Distributary complex of the Atchafalaya River; **Hma**, Distributary complex occupied by the Mississippi and Atchafalaya Rivers; **Hua**, Holocene undifferentiated alluvium).

QUATERNARY SYSTEM

HOLOCENE

Hua Holocene undifferentiated alluvium

PLEISTOCENE

LOESS
[pattern] Peoria Loess

PRAIRIE ALLOGROUP
Ppav Avoyelles alloformation
Ppbe Beaumont Alloformation

3. Units mapped in the Lafayette 7.5-minute quadrangle.



4. Correlation of strata mapped in the Lafayette 7.5-minute quadrangle.

Allostratigraphic Approach to Pleistocene Unit Definitions

In the late 1980s the LGS had begun exploring the application of allostratigraphic concepts and nomenclature to the mapping of surface Plio–Pleistocene units (e.g., Autin, 1988). In Louisiana these units show a series of geomorphic attributes and preservation states correlative with their relative ages, which eventually led LGS to conclude that allostratigraphy offers an effective if not essential approach to their delineation and classification (McCulloh et al., 2003; McCulloh, 2013). The Plio–Pleistocene strata for which allostratigraphic nomenclature presently has value to LGS all are situated updip of the hinge zone of northern Gulf basin subsidence, and show a clear spectrum of preservation from pristine younger strata to trace relicts and remnants of older strata persisting in the coastal outcrop belt and on high ridgetops in places updip of it. Allounit nomenclature has figured heavily in the STATEMAP-funded geologic mapping projects of the past two decades because Quaternary strata occupy approximately three-fourths

Lafayette Parish directly to the north (U.S. Geological Survey, 2011); the Avoyelles alloformation is sand-rich near the surface and may have potential to become a sand resource. The 1:24,000-scale surface-geologic map of the study area also should serve efforts at protection of the Chicot aquifer in the greater Lafayette area.

Acknowledgments

The work described and summarized herein was supported by the National Cooperative Geologic Mapping Program, STATEMAP component, under cooperative agreement G16AC00183 with the U.S. Geological Survey.

References

Autin, W. J., 1996, Pleistocene stratigraphy in the southern Lower Mississippi Valley:

Heinrich, P. V., J. Snead, and R. P. McCulloh (compilers), 2003, Crowley 30 × 60 minute geologic quadrangle: Louisiana Geological Survey, Baton Rouge, Scale 1:100,000.

Heinrich, P. V., J. Snead, and R. P. McCulloh (compilers), 1997, Crowley, Louisiana 30 × 60 minute geologic quadrangle (preliminary): Unpublished map prepared in cooperation with U. S. Geological Survey, STATEMAP program, under cooperative agreement no. 1434–HQ–96–AG–01490, Louisiana Geological Survey, Baton Rouge, 1:100,000-scale map plus explanation and notes.

Howe, H. V., and C. K. Moresi, 1933, Geology of Lafayette and St. Martin Parishes: Louisiana Department of Conservation, Louisiana Geological Survey, Geological bulletin no. 3, 238 p.

Milner, L. R., and C. Fisher, 2009, Geological characterization of the Chicot/Atchafalaya aquifer region: southwest Louisiana: Louisiana Geological Survey, Water resources series no. 4, 39 p.

Rouly, K. C., 1989, Quaternary and environmental geology of south-western St. Martin Parish, Louisiana: M.S. thesis, University of Southwestern Louisiana, Lafayette, Louisiana.

Saucier, R. T., 1994a, Geomorphology and Quaternary geologic history of the Lower Mississippi Valley: volume 1, Vicksburg, Mississippi, U. S. Army Corps of Engineers, Waterways Experiment Station, 364 p. plus appendices.

Saucier, R. T., 1994b, Geomorphology and Quaternary geologic history of the Lower Mississippi Valley: volume 2, Vicksburg, Mississippi, U. S. Army Corps of Engineers, Waterways Experiment Station [unpaginated: 31 oversized pages, including 28 plates (1:250,000-scale)].

Saucier, R. T., and J. I. Snead (compilers), 1989, Quaternary geology of the Lower Mississippi Valley, *in* Morrison, R. B., ed., Quaternary non-glacial geology: conterminous United