the veterinarian’s role in

Honey bees weren’t on veterinarians’ radars until ʌ t ʊ o aת q

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

-3(N)]TJET EMC /P &MCID 2 BDC BT1 0 0256 610501.43.86 Td(veepth-)3d)-ode (7)o(O)n5ag(OOd6(e)39-c5(o)ve(3(o((O)n5O)s)(tg

All honey bees in the United States are the same species, the European/Western honey bee (*Apis mellifera*). Honey bees operate as a super organism (colony), made up of individual organisms (bees). All of the functions of the colony occur within a cavity like hive, on multiple sheets of comb, which is made up of a series of . Within the colony, there are three types of honey bees based on their function:

Each colony generally contains one queen. The queen is the fertile female of the hive and is the sole source of fertilized eggs that become worker bees, laying up to 2,000 eggs per day during peak production. The queen is the largest bee in the hive, approximately twice the length of a worker bee, and longer than the drones, with a more tapered abdomen. A colony will only produce new queens when it prepares to reproductively split by swarming, when the old queen has died, or to replace a failing queen. Many queen cells will be created, and the first one to emerge will kill the remainder, and will fight with other emerged virgin queens so that only one remains. One to two weeks after hatching the virgin queen will go on several mating flights where she will mate with 10-20 drones, storing the sperm for use over her lifetime. A beekeeper may ' ' or replace a queen to introduce a younger or better performing queen. A colony can only function normally when a queen is present and laying well. A colony is considered " " when a queen is present, and " " in the absence of a queen.

Drones are the only male bees in the hive and are haploid (having only one chromosome set) because they arise from unfertilized eggs (queens and workers are diploid because they arise from fertilized eggs). Drones have large, thick bodies – larger than those of the worker bees, but shorter and thicker than that of the queen. Drones perform no functions inside the hive; their sole duty is to search for and mate with virgin queen bees on their mating flights. These flights occur far from their hives. If a drone is fortunate enough to mate, his endophallus is removed in the process and the drone dies. Drones are made whenever the colony has sufficient resources; in the summer, a strong colony can have hundreds of drones, but they are all kicked out of the colony before the winter so they don't consume precious resources.

Workers are female bees that perform the vital work of the colony. Worker bees have a variety of functions such as providing for the queen's needs; cleaning cells in the comb, nursing larvae, producing wax and forming it into honey comb, guarding and defending the hive, removing dead bees from the hive, cooling the hive or heating the brood, carrying water, gathering and transporting pollen, gathering propolis and using it to coat the hive, collecting nectar, sealing honey, and scouting for resources. Worker bees are incapable of laying fertilized

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

eggs that can become queens or other worker bees; they are only capable of laying unfertilized eggs that become drones, but this ability to lay even unfertilized eggs is suppressed in the presence of a laying queen. The lifespan of a worker bee varies with the time of year: they may live only 5-6 weeks during the spring and summer, and may live five months or longer during the inactive winter period.

Image 1: Side-by-side comparison of a worker bee, queen, and drone.

The term refers to the young, developing bees - the , and Bees at this stage remain stationary within the cells of the comb. All bees begin their lives as eggs laid in cells in the colony. After 3 days the egg hatches, and a larva emerges. The larvae are fed and grow over the next 6 days, after which the cell containing the larva is capped (the open top is sealed over by worker bees with porous wax). The larva then matures to a pupa inside the capped cell (*capped brood)* and eventually emerges from the cell as a bee. The total time spent as brood is 16 days for queens, 21 for workers, and 24 days for drones. A colony will contain brood during most of the year. Egg laying ceases in late fall or early winter and in times of stress.

The diet of honey bees is entirely from flowers. Floral nectar provides a source of carbohydrates. Bees prefer to eat fresh nectar when available, but they will store it in cells for when there are no available flowers. To prevent fermentation, the bees dry the nectar to below 18% water content, which then is referred to as honey. Pollen provides a source of protein, vitamins, fats, and minerals. To store pollen, bees will pack it into cells, add nectar, and ferment the pollen into a storable substance called bee bread.

A typical honey bee colony consists of one queen, hundreds of drones (in summer), and tens of thousands of workers. A healthy colony is one where there the queen is laying enough eggs and the workers can raise enough brood to replace the workers that are dying, and there are enough members of each age of worker to perform all the necessary tasks of the colony.

Back to Table of Contents Next: Beekeeping Equipment and Terminology

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the

Image 4:

All hives must have removable frames to allow for inspection. In image 5 below, one has been removed from an 8-frame super (hive box); the is the central portion framed by the wooden frame. The foundation is a flat piece of beeswax or plastic with the outline of cells embossed into it. The bees will create comb using the foundation as a base, a process beekeepers call ‘



Image 5: Unused frame with foundation. Photo © Jason Morgan, allmorgan.com

Image 6: Frame inspection. Photo © Randy Oliver, ScientificBeekeeping.com

The main tool used by beekeepers is the (image 7), a pry bar used to pry open the lid, separate boxes, and separate and frames for removal. A soft (image 8) may be used to gently brush bees from the removed frame to allow examination of the underlying brood. A is used to move bees away from areas to be examined and to calm the bees.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Image 7:The hive tool (yellow) has been used to safely separate frames so an individual frame can be removed without harming the bees. Photo © Randy Oliver, ScientificBeekeeping.com

Image 8: Bee brush in use to gently remove bees from frame to facilitate visual inspection. Photo © Randy Oliver, ScientificBeekeeping.com

Protection worn by beekeepers is variable, and may include coveralls (secured at wrists and leg hem); gloves; and a screened head cover with veil. Some experienced beekeepers don't wear

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

When working with a beekeeper, use only their equipment to prevent the transmission of disease between apiaries. The beekeeper should provide a smoker and hive tool, though you should have your own protective gear. Wear disposable nitrile gloves as these will prevent most stings and can be disposed of after use. When you are finished inspecting an apiary, make sure

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the

Colony health is strongly associated with its nutritional status, which is predominately a function of the amount and quality of incoming pollen. Check to see whether there is a band of stored pollen in an arc above and to the sides of the brood (image 14). A 1" arc of bee bread of multiple colors is optimal; a narrower arc, or bee bread of only one color, may indicate a food dearth. A colony should have excess stored nectar/honey and pollen at all times.

The easiest and most direct assessment of nutritional status is to see how much jelly the workers are placing around young larvae.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Queen cells are immediately recognizable because of their size and perpendicular orientation to the ground (they point towards the earth, while worker and drone brood lie horizontally/ parallel to the ground). Queen cells may indicate that there is an issue with the current queen; that the queen was killed and is being replaced; that the colony is large/crowded and will soon swarm; or they may indicate nothing – sometimes the bees make them, and later tear them down. Queen cups, which look like wax bowls facing down, are not the same as queen cells. Cups are present in the hive throughout the year and are the base for queen cells.

Image 17: Queen cells. Note the downward-pointing, large cells that are often said to resemble peanut shells.
P

Back to Table of Contents Next: Honey Bee Diseases and Conditions

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Honey bees can be host to a variety of bacterial, viral, fungal, and microsporidial diseases. Only two honey bee diseases, however, are commonly treated with antibiotics, therefore requiring veterinary oversight: European foulbrood (EFB) and American foulbrood (AFB). It is helpful for veterinarians to have a basic understanding of common honey bee diseases, and to recognize them, even if antibiotics are not indicated. It is also important to keep in mind that a colony can have multiple diseases at a time, so identification of key symptoms is vital. With all diseases, it is important to identify how heavily infected a colony is, and if the condition is worsening or getting better. To record disease severity, record the percentage of the brood demonstrating signs of disease. As with any animal, it is common for a colony to have a slight infection that it clears on its own, and slight infections of some diseases are not necessarily cause for concern.

Bacterial Diseases
- American foulbrood (AFB)
  - Visual inspection findings
  - Field tests
  - Laboratory tests and reporting
  - Treatment
- European foulbrood

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

American and European foulbrood are two significant bacterial honey bee diseases that may require veterinary intervention. Both diseases have worldwide distribution, and are commonly treated with antibiotics. The “foulbrood” name originated due to the foul smell arising from the decay of the infected brood, but American and European foulbrood are not closely related. Because these are the two known bacterial infections, these are likely the only two diseases that currently require consultation from a veterinarian.

American foulbrood (AFB)

Visual inspection findings

European foulbrood (EFB)

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been

The odor of AFB is distinctive—something between decomposing adult bees and old gym socks. European foulbrood (EFB) is either odorless or produces a sour milk-like smell. Some other unidentified diseases can also have strong smells, but none the same as AFB. A beekeeper with a good nose can easily smell AFB a few feet away from the colony entrance. A colony with a small infection may not have a noticeable odor.

The first sign consistently observed with any disease affecting the brood is a *spotty (or "shotgun") pattern* of the brood cells. This is indicative of brood disease, but is not pathognomonic for foulbrood. This indicates that brood are dying before they are capped.

Image 18: A "shotgun" brood pattern typical of American foulbrood. This frame came from a colony with an advanced case of the disease. Photo © Randy Oliver, ScientificBeekeeping.com

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Image 21: Inspecting a frame for AFB scale, looking at the bottom wall of the cell by holding the frame with the top bar towards you. Photo © Randy Oliver, ScientificBeekeeping.com

Image 22: Typical AFB infection. Note the black, raised scales lining the bottom wall of several cells, as well as the sunken, perforate caps. Photo © Randy Oliver, ScientificBeekeeping.com

– AFB kills young bees at a specific developmental stage, and they sometimes die in a characteristic manner, with the developing proboscis exposed, referred to as the 'pupal tongue', Image 23. The presence of *pupal tongues* is characteristic of AFB, but the absence of pupal tongues does not rule out the disease.



Image 23: Deceased pupae demonstrating characteristic pupal tongue
Image source: The Management Agency, National American Foulbrood Pest Management Plan, New Zealand)

Back to Table of Contents Back to Diseases Section Back to AFB main section

Next: AFB Diagnosis, Field Tests

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

A *positive rope test* (also called matchstick test) is characteristic of AFB. To perform the test, insert a matchstick, toothpick or similar object into a cell with a discolored, oozing cell cap and then slowly pull it out. The decaying products in the cell will form a viscous string from the tool, and will rope out 2 cm or more, as shown in Images 24 and 25

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Shotgun/patchy brood pattern
Dying pre-pupae or pupae melting against cell wall

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

.304 4 27 reW*3A11(931.04 P EMC P M MCID 11>>MCID 15>>BDC q93.384 15. 0.eW*nBTF1B(s)4(wn lig)5W*ni)4(n)t3(bbnBtt(fo)45e)3(d)57 ID 12>>BDC 136.58 244.13 226.25 13.56 ref*EMC P M

Samples should be sent to the USDA Agricultural Research Service (USDA-ARS) laboratory in Beltsville, Maryland.

How to send brood samples:

Comb sample should be at least 2 x 2 inches and contain as much of the dead or discolored brood as possible. NO HONEY SHOULD BE PRESENT IN THE SAMPLE.

The comb can be sent in a paper bag or loosely wrapped in a paper towel, newspaper, etc. and sent in a heavy cardboard box. AVOID wrappings such as plastic, aluminum foil, waxed paper, tin, glass, etc. because they promote decomposition and the growth of mold.

If a comb cannot be sent, the probe used to examine a diseased larva in the cell may

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

the planned burn; do not burn near buildings, vehicles, or flammable/combustible materials; do not burn on a windy day; and take other measures as needed for your individual situation. You may be directed to cover the ashes completely with soil. Check with the authorities in your area to ensure proper destruction and site management.

*Image 28: A colony being burned and buried after a positive AFB diagnosis. Because of the severity of this disease, this is the required response in some states. Irernositi <97 123.62 reW*n 17t e13ti 0 5.*

Back to Table of Contents Back to Diseases Section Back to AFB main section

Next: Honey bee drug flowchart

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

European foulbrood (EFB) is caused by *Melissococcus pluton*, a non-spore-forming bacteria, but infection is associated with a variety of bacterial strains. EFB only affects the honey bee larval stage and is more contagious than American foulbrood (AFB). It more commonly affects stressed colonies, and may resolve spontaneously if stress is reduced and honey bee health is improved. It is less severe than AFB, but can still cause devastating brood loss. In recent years, EFB has shifted its pathogenicity in the U.S.; it no longer spontaneously clears with the advent of a good nectar flow, and EFB infection is likely to persist in the hive.

Visual inspection findings
Field and laboratory testing
Treatment

Similar to AFB and other brood diseases, brood infected with EFB will exhibit a spotty brood pattern as larvae perish, and the healthy pattern of brood is disrupted.

EFB tends to only affect older larvae, and does not affect younger larvae or capped pupae. EFB-affected larvae exhibit discoloration (turn yellow or gray), and lose their normal glistening appearance. The trachea of the developing larvae often become apparent, looking like an internal skeleton of the larvae (upper left corner of image 29). EFB is often found with a variety of other bacteria so it is common to see twisted, yellow, and gray larvae together in the same frame. Drone brood may be the first to show signs of EFB, so make sure to inspect drone larvae if present.

Infected larvae often assume a twisted or corkscrew position as they die. Sometimes the larvae have a melted appearance as they die and decay, or may look deflated or malnourished when infected.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of

Oxytetracycline and tylosin have both been used to treat European foulbrood (EFB), however, *oxytetracycline is the only drug FDA approved for EFB*. Furthermore, many beekeepers feel that tylosin is not effective, and most chose to use oxytetracycline. Beekeepers also report that EFB disease may resolve without treatment, although this seems to occur less often than previously reported. Stress reduction, including feeding, and re-queening the colony, may aid this process. If the infection is not severe (less than 10% of the brood is infected), the beekeeper may choose to employ watchful waiting, where the infected frames are marked and the colony is re-inspected in a week or so to identify if the infection is spreading or improving.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Antibiotics should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow. Due to the potential for residues in honey, it is recommended that tylosin only be used in the fall and not in the spring. Furthermore, tylosin should only be used according to directions as a powdered sugar mixture immediately after mixing. Honey stored during medication periods in combs for surplus honey should be removed following final medication of the bee colony and must not be used for human food. Honey from bee colonies likely to be infected with foulbrood should not be feeding other colonies since it may be contaminated with spore

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Two brood diseases may present similarly to European foulbrood (EFB) and American foulbrood (AFB), but may not require antibiotics; these two diseases are Idiopathic Brood Disease (IBD) and Parasitic Mite Syndrome (PMS) The specific etiologic agents of these two diseases remain unknown, although it is thought that they are caused by multiple viruses, and could be found in the presence of secondary bacterial infections.

Idiopathic Brood Disease (IBD)
- Visual inspection findings
- Diagnosis
- Treatment

Parasitic Mite Syndrome (PMS)
- Inspection and diagnosis
- Treatment

Back to Table of Contents Back to Disease Section

Next: Idiopathic Brood Disease

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Associ2 1on 2017. This inform2 1on has not been approved by the AVMA Board of D1rectors or the Hous

Shotgun/patchy brood pattern
Larvae may turn yellow but retain 'C' position
Larval death in propupal stage – sticking straight up parallel to the cell
Larvae appear melted and gummy
Larvae eventually melt into dark gray
Perforated cappings
May form a loose scale

Many beekeepers report an improvement in colony health after a course of antibiotics (usually oxytetracycline or tylosin). Others report no improvement with antibiotic treatment, and others have reported that signs spontaneously clear. Many beekeepers have reported an association of IBD and exposure to some insecticides, namely fungicides.

While some reports indicate improvement of symptoms with antibiotic treatment, spontaneous clearing also occurs with IBD. Removal of diseased frames, breaking the brood cycle (e.g. by using various methods to prevent the queen from laying eggs for several weeks), and re-queening have been shown to be useful in clearing up infection.

Back to Table of Contents Back to Diseases Section Next: Parasitic Mite Syndrome

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Varroa mites (*Varroa destructor*) are ectoparasites with worldwide distribution (reported in the U.S. since 1987) and remain the number one killer of honey bees. According to the USDA, beekeepers have identified varroa mites as their single most serious problem causing colony losses today. The mites feed on the hemolymph and fat cells of the bees. They target larvae that are about to be capped, and move to the bottom of the cell and feed off of the larva once the cell is capped. The mites mate inside the cell and mature; once the bee emerges from the cell, it will already have female mite offspring on it. The mites can cause larval or pupal death and can transmit deformed wing virus, acute bee paralysis virus, Israeli acute paralysis virus, slow bee paralysis virus and other pathogens. Affected larvae may spiral up the cell, eventually curling around the opening of the cell in a half-moon shape.

Parasitic mite syndrome (PMS) is caused by viruses transmitted by the varroa mite. In a severely mite-infested colony, bees may express parasitic mite syndrome This syndrome is most commonly seen in the

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Because the mites carry multiple viruses, and because a

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Often, young bees will be present exhibiting wrinkled, deformed wings. They may be seen moving on the frames, or crawling outside of the hive. You may not see any bees with deformed wings as they often abandon the hive or are removed.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.



© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Parasitic mite syndrome can only be prevented and controlled by managing the populations of varroa mites. Depending on the time of year and the location, it may not be possible to save a colony that is showing signs of PMS. Beekeepers in warmer climates can sometimes have a colony rebound with treatment and feeding, while colonies exhibiting PMS before a winter rarely survive. It is recommended to apply a treatment of a miticide to kill the mites and prevent the spread to other colonies.

Varroa mites are found in every hive in the country and are the most devastating parasites of bees because of their ability to vector other disease agents. Beekeepers must have a strategy in place to manage the varroa mite, and those that do not will experience high mortality of their hives and can transmit disease to nearby colonies. Mite-associated disease remains one of the leading causes of honey bee death, especially among hobby and small scale beekeepers. The honey bee veterinarian should know about the life cycle of this parasite, its monitoring methods, control points, and treatment methods. See the following resources for more information:

Honey bee health coalition tools to manage varroa
Michigan State varroa resources

Small- scale beekeepers should be practicing integrated pest management (IPM) to manage for varroa mites. This means that they are actively monitoring mite populations (generally with an alcohol wash or sugar roll), and should employ many physical, mechanical, and if necessary chemical controls. Most commercial beekeepers are able to manage mite populations through

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Two types of Nosema infections are possible, caused by the single-celled microsporidia *Nosema apis* and *Nosema ceranae*. The spores spread from bee to bee by feeding and fecal-oral transmission. The two Nosema diseases have different natural histories. *Nosema apis* tends to be worse in northern climates, and outbreaks occur more often in the fall and winter when bees cluster together to maintain hive warmth, while *Nosema ceranae* can be present year round.

Signs of Nosema disease can include bees wandering on ground outside colony; "K-wing," (see Image 45), which is an uncoupling of the forward and rear pairs of wings that results in wings that resemble the letter K; yellow, orange or brown fecal staining on front of colony; and poor growth of bees and death.

. Nosema infection must be confirmed by simple microscopy, either of a sample of adults, or by one-by-one gut squashes.

Image 40 - Nosema spores at 400x in a hemacytometer. Photo © Randy Oliver, ScientificBeekeeping.com

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Chalkbrood is caused by the *Ascosphaera apis* fungus that attacks larvae and pupae. This condition is commonly seen in the spring and is more prevalent during rainy or wet weather, or after the colony has been chilled.

White fungal growth is observed, followed by white or black "mummies" of dead larvae or pupae. These mummies are seen in the brood cells, and also at the entrance of the hive.

There is no chemical therapy for chalkbrood. Stressed hives are at higher risk, and the disease is best controlled by reducing stress through good management practices and by using disease-resistant bees. The hygienic behavior of a

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Tracheal mite infestation
Small hive beetles
Tropilaelaps spp

Tracheal mite infestation (Acarapisosis) can be difficult to recognize until the damage is severe. Mite females infest 1-2 day old bees and penetrate their thorax and trachea and lay eggs. The offspring mature and mate inside the trachea, causing damage obstruction. Crowded colonies are at higher risk, and infestations tend to peak in the fall and winter when bees are clustered. Affected bees may develop "K-wing," (Image 45) which is an uncoupling of the forward and rear pairs of wings that results in wings that resemble the letter K; however, this is not unique to tracheal mite infestation. Confirmation of infestation is made with microscopic evaluation of affected bees.

This disease has not been widely found in the USA in recent

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

Tropilaelaps is the genus name of a mite with four species identified. None have been found in the U.S. When *Apis melifera* has been kept in regions where this mite is found, the mite can create severe infection. Tropilaelaps causes damage similar to varroa, resulting in irregular brood patterns and stunted adults with deformed wings and shrunken abdomens. Tropilaelaps mites are reddish brown, and are smaller than the varroa mite - about 1mm long and 0.6mm wide - and can be seen running on frames in between cells. Tropilaelaps is of high regulatory concern and any suspicious infections should be sampled and the state bee inspector contacted immediately.

Back to Table of Contents Back to Disease Section Next: Other Health Conditions

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

laying worker colonies can be shaken out and the equipment removed (the bees will enter other hives). They can also be directly added to other colonies in the yard. Large laying worker colonies should be combined with a nuc or other colony using a newspaper combine.

In 2006, beekeepers began observing dramatic declines in honey bee colonies. Further investigation revealed that the situation was different from previously known causes of loss. Colony Collapse Disorder is defined as a dead colony with no adult bees and no dead bee bodies but with a live queen, and usually honey and immature bees (brood), still present. It does not include honey bee colonies that are lost due to any reason, such as losses due to poor management or nutrition; varroa mites; or other diseases described in this section, even though the colony size may seem to 'collapse' as in the case with varroa/virus infections.

Although intensive research continues and numerous rumors and theories have been suggested, the cause for CCD has not yet been proven. According to the USDA, most research has pointed to a complex of factors being involved in the cause of CCD, though these are not uniform in all CCD incidents.

As research continues and more is learned about CCD, the USDA recommends that beekeepers focus on improving general honey bee health and habitat and use best management practices to reduce stress and mitigate the known causes of reduced production and honey bee death. An emergency response kit is available to evaluate a collapsing hive and protocols are available for sampling live bee and salt water solution and pollen.

Back to Table of Contents Back to Disease Section Next: Specimen Submission

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

The following instructions have been excerpted from the United States Department of Agriculture-Agricultural Research Service (USDA-ARS) Bee Disease Diagnosis Service site.

Beekeepers, bee businesses, and regulatory officials may submit samples.
Samples are accepted from the United States and its territories; samples are NOT accepted from other countries.
Include a short description of the problem along with your name, address, phone number or e-mail address.
There is no charge for this service.
For additional information, contact Sam Abban by phone at (301) 504-8821 or e-mail at samuel.abban@ars.usda.gov.

Send at least 100 bees and if possible, select bees that are dying or that died recently.
Decayed bees are not satisfactory for examination.
Bees should be placed in and soaked with 70% ethyl, methyl, or isopropyl alcohol as soon as possible after collection and packed in leak-proof containers.
USPS, UPS, and FedEx do not accept shipments containing alcohol. Just prior to mailing samples, pour off all excess alcohol to meet shipping requirements.
Do NOT send bees dry (without alcohol).

A comb s Tm[si)12(te1f3(s)]TJET EMC /P <MCIDIul1 0 -3(wi)ed w)(l).) be 24(o)xJET ETC c(e)5()10(in l)13(TJET

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of

USDA: Honey Bees
Honey bee lifecycle (South Carolina Mid-State Beekeepers Association)

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

(video) European and American foulbrood in honey bee colonies: Part 1 (University of Florida Honey Bee Research & Extension Lab)
(video) European and American foulbrood in honey bee colonies: Part 2 (University of Florida Honey Bee Research & Extension Lab)
(video) Identifying American Foulbrood (DPI Agriculture, Australia)
(video) American Foulbrood (Crispin Boxhall)
(video) AFB elimination 8 – how to recognise AFB (New Zealand)
(video) European foulbrood (Crispin Boxhall)

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.

The AVMA would like to recognize the important contributions of the following veterinarians and expert beekeepers to this guide. Without their input, this comprehensive reference would not have been possible.

Dr. Chris Cripps
Dr. Christine Hoang
Dr. Don Hoenig
Dr. Kimberly May
Dr. William McBeth
Dr. Meghan Milbrath
Randy Oliver
Dr. Robert Wills
Dr. Jennifer Wishnie

Back to Table of Contents

© American Veterinary Medical Association 2017. This information has not been approved by the AVMA Board of Directors or the House of Delegates, and it is not to be construed as AVMA policy nor as a definitive statement on the subject, but rather to serve as a resource providing practical information for veterinarians.