Office of Research and Graduate Education

Sample abstracts are enclosed for additional information.

Template abstract 1 - Research

Differentiation of Canine Induced Pluripotent Stem Cells into Neural Progenitor Cells

XXXX
Laboratory for XXX, Department of XXX, XXX University

Background and Rationale: New advances in stem cell technology, including induced pluripotent stem cells (iPSC), offers new hope for patients with neurological disease and spinal cord injuries. Therefore, we evaluated the ability of canine iPSC to be differentiated into neural progenitor cells (NPC) in vitro as a precursor to clinical trials in dogs with spinal cord injury.

Approach: iPSC were generated from canine fibroblasts and characterized based on phenotype, gene expression analysis, lineage differentiation, and teratoma formation. Canine iPSC were then induced to differentiate into NPC by culture in defined medium supplemented with specific growth factors. NPC were characterized by phenotype, flow cytometry, immunofluorescence, and gene expression analysis.

Results: Canine iPSC could be readily induced to differentiate into NPC following 2-3 weeks in culture. Specific culture conditions led to enrichment of NPC for cells with characteristics of oligodendrocytes, astrocytes and neurons. NPC did not form teratomas in mice, whereas the parental iPSC cells did.

Conclusions/Implications: Canine iPSC can be induced to form NPC in vitro by altering cell culture conditions, cell substrate, and addition of specific growth factors. These studies provide evidence that iPSC technology can be used to generate NPC for use in neural regeneration in dogs with neurological injuries.

Template abstract 2 - Research

Key Role For Scavenger Receptor B-I In The Integrative Physiology Of Host Defense During Bacterial Pneumonia

K.M. XXX, M.B. XXX,
Department of XXXXXXXXXXXXXX

Rationale: Scavenger Receptor B-I (SR-BI) is a multi-recognition receptor mostly studied in the arena of atherosclerosis due to its role in cellular uptake of cholesterol ester from high density lipoprotein (HDL). Recently, SR-BI has also been reported to play a role in clearance of lipopolysaccharide (LPS) from the plasma.

Methods: SR-BI+/+ and SR-BI-/- mice were infected intratracheally with *Klebsiella pneumoniae* (Kp), or challenged with aerosolized LPS. Survival, leukocyte influx into the airspace, bronchoalveolar lavage fluid (BALF) cytokines, and bacterial CFUs in lung and blood were quantified. Bactericidal function of neutrophils (PMNs) was evaluated.

Results: Compared to SR-BI+/+ counterparts, SR-BI-/- mice suffered markedly increased mortality during pneumonia, in conjunction with higher bacterial burden in lung and blood, deficient induction in the plasma of the stress glucocorticoid corticosterone, higher serum cytokines, and increased peripheral organ injury. SR-BI-/- mice had significantly enhanced PMN recruitment to the airspace as well as increased BALF TNF-α, G-CSF, and CXCL5 after pulmonary exposure to either Kp or LPS. This was associated with defective clearance of LPS from the SR-BI-/- airway and increased cytokine production by SR-BI-/- macrophages. SR-BI-/- PMNs displayed decreased phagocytosis and a dramatic defect in intracellular bacterial killing.

Conclusions: SR-BI is central regulator of the integrated pulmonary host defense response through multiple interacting mechanisms, including modulation of alveolar cytokine induction, PMN recruitment, and PMN function.

[Template abstract 3 - Research]

Pharmacokinetics and Pharmacodynamics of an Extended Release Buprenorphine Formulation in Dogs

XXXX
Deparep-4.-dm78aβ13 0.205 0 :,

Template abstract 5 – Case report

Extramedullary Plasmacytoma in a Hamster

X. XXX, X.X XXX
Department of XXXXXXXXX

Background: An extramedullary plasmacytoma is a neoplastic proliferation of plasma cells arising from outside of the bone marrow. It is a relatively common tumor in old dogs and occurs most frequently on the skin and mucous membranes. Salivary gland plasmacytomas are extremely rare within all species.

Case Description: A 1.5-year-old, intact female Syrian hamster (*Mesocricetus auratus*) showed respiratory distress and a ventral cervical mass was palpated. Due to the overall poor prognosis, the animal was humanely euthanized and submitted for necropsy. On gross examination a subcutaneous, round, loosely attached, semi-firm mass measuring approximately 1 cm in diameter was identified at the ventral neck region. Histopathologic examination of the cervical mass revealed neoplastic proliferation of mononuclear cells with plasmacytoid morphology adjacent to and infiltrating the submaxillary salivary gland. The neoplastic cells were positive for CD79a and negative for CD3. Based on the diagnostic test results and histopathologic evaluation of the mass tissue sections, a diagnosis of an extramedullary plasmacytoma was

Conclusion: Pulmonary toxoplasmosis is generally a diffuse, interstitial disease, whereas this dog had an unusual, multi-nodular, pyogranulomatous presentation.